## NIDA Quick Screen V1.01

Interviewer...... Date ...../.....

| Introduction (Please read to patient)  |           |       |                  |         |        |                             |
|--|-----------|-------|------------------|---------|--------|-----------------------------|
| Hi, I'm, nice to meet you. If it's okay with you, I'd like to ask you a few questions that will help me give you better medical care. The questions relate to your experience with alcohol, cigarettes, and other drugs. Some of the substances we'll talk about are prescribed by a doctor (like pain medications). But I will only record those if you have taken them for reasons or in doses other than prescribed. I'll also ask you about illicit or illegal drug use—but only to better diagnose and treat you.  Instructions: For each substance, mark in the appropriate column. For example, if the patient has used cocaine monthly in the past year, put a mark in the "Monthly" column in the "illegal drug" row. |           |       |                  |         |        |                             |
| NIDA <i>Quick Screen</i> Question:  In the past year, how often have you used the fe   | ollowing? | Never | Once or<br>Twice | Monthly | Weekly | Daily or<br>Almost<br>Daily |
| Alcohol  |           |       |                  |         |        |                             |
| <ul> <li>For men, 5 or more drinks a day</li> <li>For women, 4 or more drinks a day</li> </ul>   |           |       |                  |         |        |                             |
| Tobacco Products   |           |       |                  |         |        |                             |
| Prescription Drugs for Non-Medical Reasons   |           |       |                  |         |        |                             |
| Illegal Drugs  |           |       |                  |         |        |                             |

- If the patient says "NO" for all drugs in the Quick Screen, reinforce abstinence. Screening is complete.
- If the patient says "Yes" to one or more days of heavy drinking, patient is an at-risk drinker. Please see NIAAA website "How to Help Patients Who Drink Too Much: A Clinical Approach" <a href="http://pubs.niaaa.nih.gov/publications/Practitioner/CliniciansGuide2005/clinicians guide.htm">http://pubs.niaaa.nih.gov/publications/Practitioner/CliniciansGuide2005/clinicians guide.htm</a>, for information to Assess, Advise, Assist, and Arrange help for at risk drinkers or patients with alcohol use disorders
- If patient says "Yes" to use of tobacco: Any current tobacco use places a patient at risk. Advise all tobacco users to quit. For more information on smoking cessation, please see "Helping Smokers Quit: A Guide for Clinicians" <a href="http://www.ahrq.gov/clinic/tobacco/clinhlpsmksqt.htm">http://www.ahrq.gov/clinic/tobacco/clinhlpsmksqt.htm</a>
- If the patient says "Yes" to use of illegal drugs or prescription drugs for non-medical reasons, proceed to Question 1 of the NIDA-Modified ASSIST.

This guide is designed to assist clinicians serving adult patients in screening for drug use. The NIDA Quick Screen was adapted from the single-question screen for drug use in primary care by Saitz et al. (available at <a href="http://archinte.ama-assn.org/cgi/reprint/170/13/1155">http://archinte.ama-assn.org/cgi/reprint/170/13/1155</a>) and the National Institute on Alcohol Abuse and Alcoholism's screening question on heavy drinking days (available at <a href="http://pubs.niaaa.nih.gov/publications/Practitioner/CliniciansGuide2005/clinicians\_guide.htm">http://pubs.niaaa.nih.gov/publications/Practitioner/CliniciansGuide2005/clinicians\_guide.htm</a>). The NIDA-modified ASSIST was adapted from the World Health Organization (WHO) Alcohol, Smoking and Substance Involvement Screening Test (ASSIST), Version 3.0, developed and published by WHO (available at <a href="http://www.who.int/substance\_abuse/activities/assist\_v3\_english.pdf">http://www.who.int/substance\_abuse/activities/assist\_v3\_english.pdf</a>).